## GETTING THE BEST FROM ACELEPRYN



## YOUR 7 POINT PLAN FOR APPLICATION



Apply preventatively. Acelepryn is only effective on 1st and 2nd instar. Larger grubs / Larvae will not be controlled.



Apply at 0. 6L/ha, in a minimum of 600 litres of water, using a Syngenta XC soil nozzle or suitable air induction 08 nozzle between 2–4 bar pressure.



Chafers: Apply once peak flight has been observed (typically June).

LeatherJackets: Apply one month post peak flight (UK peak flight typically Mid September).

Sites with historic high pressure have increased probability of re-infestation.

Report sightings to PestTracker at:

www.syngentaturf.co.uk/turf-pest-tracker



Ensure the soil profile is not hydrophobic at the time of application. Syngenta wetting agent Qualibra is compatible with Acelepryn for convenient application in tank-mix or sequence.



Keep surfaces irrigated where possible, to attract pest larvae to towards the surface.





Delay mowing for as long as possible and return the clippings if feasible.

## RESEARCH HAS SHOWN ACELEPRYN PROVIDES UP TO 90% CONTROL WITH A WELL PLANNED AND EXECUTED APPLICATION

## HIGH AND MIXED POPULATIONS WILL NEED A LONG TERM PROGRAMMED APPROACH

Follow #PestTracker for timing information support and @SyngentaTurfUK for turf management advice. For technical support contact Sean Loakes on sean.loakes@syngenta.com or 07831 161972.

Syngenta UK Ltd. Registered in England No. 849037. CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE. Email: customer.services@syngenta.com.

Acelepryn® and Qualibra® are Registered Trademarks of a Syngenta Group Company. Acelepryn contains chlorantraniliprole. All other brand names used are trademarks of other manufacturers in which proprietary rights may exist. Use plant protection products safely. Always read the label and product information before use. For further product information including warning phrases and symbols refer to www.syngentaturf.co.uk ©Syngenta AG June 2022. GQ 12105.

